

List of Plants

C (E)-BETA-OCIMENE

Chemid

EBETAOCIMENE

*Unless otherwise noted all references are to Duke, James A. 1992. Handbook of phytochemical constituents of GRAS herbs and other economic plants. Boca Raton, FL. CRC Press.

Plant	Plant Part	Low PPM	High PPM	StdDev	*Reference
Apium graveolens	Seed	7.0	10.0		Charalambous, G. (Ed.). 1994. Spices, Herbs and Edible Fungi. Elsevier Science B. V. Amsterdam. 764 pp.
Carica papaya	Fruit	--	--		*
Carum carvi	Seed	--	--		*
Cinnamomum verum	Leaf	--	--	-1.14	Mallavarupu, G. R. et al. 1995. Investigation of the essential oil of cinnamon leaf grown at Bangalore and Hyderabad. Flav. & Fragr. J., 10: 239-242.
Citrus sinensis	Fruit	0.7	1.0	-0.8	*
Citrus limon	Petiole	--	6.0		*
Citrus reticulata	Fruit	4.0	5.0	-0.62	*
Citrus paradisi	Pericarp	--	8.0		*
Dictamnus albus	Shoot	--	6.0	-0.6	Baser, K.H.C., Kosar, M.Malyer, H. & Ozek, T. 1994. The Essential Oil Composition of Dictamnus albus from Turkey. Plant a Med. 60:481-2
Dictamnus albus	Shoot	--	6.0	-0.6	Baser, K.H.C., Kosar, M.Malyer, H. & Ozek, T. 1994. The Essential Oil Composition of Dictamnus albus from Turkey. Plant a Med. 60:481-2
Dictamnus albus	Shoot	--	7.0	-0.59	Baser, K.H.C., Kosar, M.Malyer, H. & Ozek, T. 1994. The Essential Oil Composition of Dictamnus albus from Turkey. Plant a Med. 60:481-2
Elsholtzia blanda	Shoot	--	38.0	-0.2	Bestman, H.J., Rauscher, J., Vostrowsky O., Pant, A.K., Dev. V., Perihar, R. and Mathela, C.S. 1992. Constituents of the Essential Oil of Elsholtzia blanda Benth. (Labiatae). J. Ess. Oils Res. 4: 121-124
Eucalyptus bridgesiana	Leaf	--	8.0	-0.61	Singh, A. K., Gupta, K. C., & Brophy, J. J. 1991. Volatile Constituents of the Essential Oil of Eucalyptus bridgesiana Growing in India. Journal of Essential Oil Res. 3: 449-450.
Hyptis suaveolens	Shoot	--	--	-0.68	Mallvarapu, G.R., Ramesh, S., Kaul, P.N., Bhattacharya, A.K., and Rao, B.R.R. 1993. The Essential Oil of Hyptis suaveolens (L.) Poit. J. Ess. Oil Res. 5: 321.
Hyptis suaveolens	Shoot	--	--	-0.68	Mallvarapu, G.R., Ramesh, S., Kaul, P.N., Bhattacharya, A.K., and Rao, B.R.R. 1993. The Essential Oil of Hyptis suaveolens (L.) Poit. J. Ess. Oil Res. 5: 321.
Hyssopus officinalis	Shoot	--	70.0	0.21	Tsankova, E.T., Konatchiev, A.N. and Genova, E.M. 1993. Chemical Composition of the Essential Oils of Two Hyssopus officinalis cultivars. J. Ess. Oil Res. 5: 609-611.
Hyssopus officinalis	Shoot	--	70.0	0.21	Tsankova, E.T., Konatchiev, A.N. and Genova, E.M. 1993. Chemical Composition of the Essential Oils of Two Hyssopus officinalis cultivars. J. Ess. Oil Res. 5: 609-611.
Lavandula x hybrida	Shoot	22.0	48.0	-0.07	Tucker, A.O., Maciarello, M.J., Angell, S., Espaillat, J.R., and French, E.C. 1993. The Essential Oil of Lavandula x hybrida Balb. ex Ging., a Distinct Hybrid from L. x heterophylla Poir. (Labiatae). J. Ess. Oil Res. 5: 443-445.
Leonotis leonurus	Se	--	3.0		Pedro, L.G., Barroso, J.G., Marques, N.T., Ascenso, L., Pais, M.S.S. and Scheffer, J.J.C. 1991. Composition of the Essential Oil from Sepals of Leonotis leonurus R. Br. J. Ess. Oil Res. 3: 451-3
Lindera benzoin	Fruit	36.0	50.0	1.41	*
Micromeria myrtifolia	Shoot	--	0.3	-0.67	Ozek, T., Kirimer, N., and Baser, K.H.C. 1992. Composition of the Essential Oil of Micromeria myrtifolia Boiss. et Hohen. J. Ess. Oil Res., 4: 79-80.
Micromeria fruticosa	Shoot	--	19.0	-0.44	Fleisher, Z. and Fleisher, A. 1991. The Essential Oil of Micromeria fruticosa (L.) Druce subsp. barbata (Boiss et. Ky.), P.H. Davis. Aromatic Plants of the Holy Land and the Sinai. Part VII. J. Ess. Oil Res 3: 477-479.
Micromeria fruticosa	Shoot	--	19.0	-0.44	Fleisher, Z. and Fleisher, A. 1991. The Essential Oil of Micromeria fruticosa (L.) Druce subsp. barbata (Boiss et. Ky.), P.H. Davis. Aromatic Plants of the Holy Land and the Sinai. Part VII. J. Ess. Oil Res 3: 477-479.
Minthostachys mollis	Shoot	3.0	18.0	-0.45	Alkire, B.H., Tucker, A.O., and Maciarello, M.J. 1994. Tipo (Minthostachys mollis (Lamiaceae): An Ecuadorian Mint. Econ. Bot. 48(1): 60-64.
Murraya koenigii	Leaf	--	40.0	1.5	*
Ocimum basilicum	Plant	1.0	435.0	2.23	*
Origanum vulgare	Plant	--	40.0	-0.37	Sezik, E., Tumen, G., Kirimer, N., Ozek, T., and Baser, K.H.C. 1993. Essential Oil Composition of Four Origanum vulgare Subspecies of Anatolian Origin. J. Ess. Oil Res., 5: 425-431.

sipyleum					sipyleum of Turkish Origin. J. Ess. Oil Res. 4: 139-142.
Origanum minutiflorum	Shoot	--	5.0	-0.61	Baser, K.H.C., Tumen, G., Sezik, E. 1991. The Essential Oil of Origanum minutiflorum O. Schwarz and P.H. Davis. J. Ess. Oil Res. 3: 445-446.
Origanum vulgare	Plant	--	25.0	-0.47	Sezik, E., Tumen, G., Kirimer, N., Ozek, T., and Baser, K.H.C. 1993. Essential Oil Composition of Four Origanum vulgare Subspecies of Anatolian Origin. J. Ess. Oil Res., 5: 425-431.
Origanum vulgare	Plant	--	25.0	-0.47	Sezik, E., Tumen, G., Kirimer, N., Ozek, T., and Baser, K.H.C. 1993. Essential Oil Composition of Four Origanum vulgare Subspecies of Anatolian Origin. J. Ess. Oil Res., 5: 425-431.
Origanum sipyleum	Shoot	--	1.8	-0.66	Baser, K.H.C., Ozek, T., Kurkcuoglu, M. and Tumen, G. 1992. Composition of the Essential Oil of Origanum sipyleum of Turkish Origin. J. Ess. Oil Res. 4: 139-142.
Origanum vulgare	Plant	--	50.0	-0.31	Sezik, E., Tumen, G., Kirimer, N., Ozek, T., and Baser, K.H.C. 1993. Essential Oil Composition of Four Origanum vulgare Subspecies of Anatolian Origin. J. Ess. Oil Res., 5: 425-431.
Origanum sipyleum	Shoot	--	17.0	-0.46	Baser, K.H.C., Ozek, T., Kurkcuoglu, M. and Tumen, G. 1992. Composition of the Essential Oil of Origanum sipyleum of Turkish Origin. J. Ess. Oil Res. 4: 139-142.
Origanum sipyleum	Shoot	--	70.0	0.21	Baser, K.H.C., Ozek, T., Kurkcuoglu, M. and Tumen, G. 1992. Composition of the Essential Oil of Origanum sipyleum of Turkish Origin. J. Ess. Oil Res. 4: 139-142.
Pelargonium citrosum	Shoot	--	1.0	-0.67	Matsuda, B. M., et al. 1996. Essential Oil Analysis and Field Evaluation of the Citrosa Plant 'Pelargonium citrosum' as a Repellent Against Populations of Aedes Mosquitoes. J. Am. Mosq. Contr. Assoc. 12(1):69-74.
Porophyllum ruderale	Plant	3.0	4.0	-0.61	Loayza, I., de Groot, W., Lorenzo, D. et al. 1999. Composition of the essential oil of Porophyllum ruderale (Jacq.) Cass. from Bolivia. Flav. & Fragr. J. 14: 393-8.
Rosmarinus officinalis	Shoot	--	380.0	4.15	Soriano Cano, M.C., Sotomayor Sanchez, J.A., Sanchez Gomez, P. and Garcia Vallejo, M.C. 1993. Essential Oils of the Rosmarinus eriocalyx-tomentosus Complex in Southeast Spain. J. Ess. Oil Res. 5: 243-246.
Rosmarinus eriocalyx	Shoot	60.0	185.0	1.67	Soriano Cano, M.C., Sotomayor Sanchez, J.A., Sanchez Gomez, P. and Garcia Vallejo, M.C. 1993. Essential Oils of the Rosmarinus eriocalyx-tomentosus Complex in Southeast Spain. J. Ess. Oil Res. 5: 243-246.
Rosmarinus x mendizabalii	Shoot	0.1	33.0	-0.26	Soriano Cano, M.C., Sotomayor Sanchez, J.A., Sanchez Gomez, P. and Garcia Vallejo, M.C. 1993. Essential Oils of the Rosmarinus eriocalyx-tomentosus Complex in Southeast Spain. J. Ess. Oil Res. 5: 243-246.
Rosmarinus officinalis	Shoot	12.0	25.0	-0.36	Soriano Cano, M.C., Sotomayor Sanchez, J.A., Sanchez Gomez, P. and Garcia Vallejo, M.C. 1993. Essential Oils of the Rosmarinus eriocalyx-tomentosus Complex in Southeast Spain. J. Ess. Oil Res. 5: 243-246.
Rosmarinus x lavandulaceus	Shoot	0.1	30.0	-0.3	Soriano Cano, M.C., Sotomayor Sanchez, J.A., Sanchez Gomez, P. and Garcia Vallejo, M.C. 1993. Essential Oils of the Rosmarinus eriocalyx-tomentosus Complex in Southeast Spain. J. Ess. Oil Res. 5: 243-246.
Rosmarinus officinalis	Shoot	--	13.0	-0.51	Soriano Cano, M.C., Sotomayor Sanchez, J.A., Sanchez Gomez, P. and Garcia Vallejo, M.C. 1993. Essential Oils of the Rosmarinus eriocalyx-tomentosus Complex in Southeast Spain. J. Ess. Oil Res. 5: 243-246.
Rosmarinus officinalis	Shoot	0.1	13.0	-0.51	Soriano Cano, M.C., Sotomayor Sanchez, J.A., Sanchez Gomez, P. and Garcia Vallejo, M.C. 1993. Essential Oils of the Rosmarinus eriocalyx-tomentosus Complex in Southeast Spain. J. Ess. Oil Res. 5: 243-246.
Rosmarinus eriocalyx	Shoot	60.0	185.0	1.67	Soriano Cano, M.C., Sotomayor Sanchez, J.A., Sanchez Gomez, P. and Garcia Vallejo, M.C. 1993. Essential Oils of the Rosmarinus eriocalyx-tomentosus Complex in Southeast Spain. J. Ess. Oil Res. 5: 243-246.
Rosmarinus tomentosus	Shoot	0.1	20.0	-0.42	Soriano Cano, M.C., Sotomayor Sanchez, J.A., Sanchez Gomez, P. and Garcia Vallejo, M.C. 1993. Essential Oils of the Rosmarinus eriocalyx-tomentosus Complex in Southeast Spain. J. Ess. Oil Res. 5: 243-246.
Sassafras albidum	Leaf	9.4	21.0	0.25	*
Satureja cilicica	Shoot	--	55.0	0.02	Tumen, G. Baser, K.H.C. and Kirimer, N. 1993. The Essential Oil of Satureja cilicica P.H. Davis. J. Ess. Oil Res. 5: 547-548.
Tagetes lucida	Shoot	--	175.0	1.55	Bicchi, C., Fresia, M., Rubiolo, P., Monti, D., Franz, C., Goehler, I. 1997. Constituents of Tagetes lucida Cav. ssp. lucida essential oil. Flavor & Fragrance, 12(1): 47-52.
Thymus zygis	Shoot	--	75.0	0.27	Jimenez, J., Navarro, M.C., Montilla, M.P., Martin, A. and Martinez, A. 1993. Thymus zygis Oil: Its Effects on CCl4-Induced Hepatotoxicity and Free Radical Scavenger Activity. JE05: 153-8.